



The  
Patent  
Office

PCT/EP 00 / 06429

REC'D 13 SEP 2000	INVENTOR PEOPLE
WIPO PCT	
The Patent Office	

Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

10/030681

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

EP 00/06429

25/01

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

E J K W

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed *Andrew G. G. G.*  
Dated 24 MAY 2000

**THIS PAGE BLANK (USPTO)**

# The Patent Office


15JUL99 EAG2169-1 D00192  
P01/7700 0.00 - 9916562.3

## Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form))

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH



1. Your reference N.77330 GCW/CMK

2. Patent application number  
(The Patent Office will fill in this part)

**9916562.3**

**14 JUL 1999**

3. Full name, address and postcode of the or of each applicant (underline all surnames)

PHARMACIA & UPJOHN S.p.A  
Via Robert Koch 1.2  
20152 Milan  
Italy

Patents ADP number (if you know it)

7/0000/001

If the applicant is a corporate body, give the country/state of its incorporation

Italy

4. Title of the invention

3-ARYLSULFONYL-2-(SUBSTITUTED-METHYL)  
PROPANOIC ACID DERIVATIVES AS MATRIX  
METALLOPROTEINASE INHIBITORS

5. Name of your agent (if you have one)

J A KEMP & CO

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

14 SOUTH SQUARE  
GRAY'S INN  
LONDON WC1R 5LX

Patents ADP number (if you know it)

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer "Yes" if:

Yes

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body:
- See note (d))

# Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 62

Claim(s) 12

Abstract 1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77) 2 x 7

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents  
(please specify)

11. I/We request the grant of a patent on the basis of this application

Signature J.A. Kemp & Co

Date 13<sup>TH</sup> July 1999

12. Name and daytime telephone number of person to contact in the United Kingdom Mrs C.M. KEEN  
0171 405 3292

## Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

## Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered "Yes" Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

# Statement of inventorship and of right to grant of a patent

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

14 JUL 1999

1. Your reference	N.77330 GCW/CMK	
2. Patent application number (If you know it)	<b>9916562.3</b>	14 JUL 1999
3. Full name of the or of each applicant	PHARMACIA & UPJOHN S.p.A	
4. Title of the invention	3-ARYLSULFONYL-2-(SUBSTITUTED-METHYL) PROPANOIC ACID DERIVATIVES AS MATRIX METALLOPROTEINASE INHIBITORS	
5. State how the applicant(s) derived the right from the inventor(s) to be granted a patent	As Employer	
6. How many, if any, additional Patents Forms 7/77 (see note (c)) are attached to this form?	1	
7.	<p>I/We believe that the person(s) named over the page (and on any extra copies of this form) is/are the inventor(s) of the invention which the above patent application relates to.</p> <p>Signature <i>JA Kemp</i> x6 Date 13<sup>TH</sup> July 1999</p>	
8. Name and daytime telephone number of person to contact in the United Kingdom	Mrs C.M. KEEN 0171 405 3292	

## Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there are more than three inventors, please write the names and addresses of the other inventors on the back of another Patents Form 7/77 and attach it to this form.
- When an application does not declare any priority, or declares priority from an earlier UK application, you must provide enough copies of this form so that the Patent Office can send one to each inventor who is not an applicant.
- Once you have filled in the form you must remember to sign and date it.

Enter the full names, addresses and postcodes of the inventors in the boxes and underline the surnames

Sergio Mantega  
Via C. Pisacane  
57-20129 Milan  
Italy

Patents ADP Number (if you know it) 557044504

Francesca Abrate  
Via Lodovico Settala  
45-20124 Milan  
Italy

Patents ADP number (if you know it) 725987202

Pierluigi Bissolino  
Via Roma  
36/2-27020 San Giorgio Lomellina  
Pavia  
Italy

Patents ADP number (if you know it) 4450034002

Reminder

Have you signed the form?

Enter the full names, addresses and postcodes of the inventors in the boxes and underline the surnames

Paolo Cremonesi  
Via San Michele del Carso  
10-20066 Melzo Milan  
Italy

Patents ADP Number (if you know it) 694228001

Ettore Perrone  
Via Aldo Moro  
44-20010 Boffalora Ticino Milan  
Italy

Patents ADP number (if you know it) 4401337001

Daniela Jabes  
Via Costanze  
33-20146 Milan  
Italy

Patents ADP number (if you know it) 59122349001

Reminder

Have you signed the form?





### 3-ARYLSULFONYL-2-(SUBSTITUTED-METHYL)PROPANOIC ACID DERIVATES AS MATRIX METALLOPROTEINASE INHIBITORS

#### FIELD OF THE INVENTION

5       The present invention relates to some  $\beta$ -sulfonyl hydroxamic and carboxylic acids displaying potent and selective inhibitory activity against some Matrix MetalloProteinases (MMPs), to their use for the treatment of diseases in which the proteolytic activity of these MMPs is involved, to methods for their preparation, and to pharmaceutical compositions containing them.

#### 10       BACKGROUND OF THE INVENTION

Under physiological conditions MMPs play an important role in normal tissue turnover and development. However, increased MMP activity and/or impairment of the physiological MMP inhibitors has been associated with a variety of pathological conditions such as cancer, rheumatoid arthritis, osteoarthritis, osteoporosis, multiple  
15   sclerosis, stroke, arteriosclerosis, restenosis and age-related macular degeneration. In cancer, the contribution of MMPs is manifold and not limited to destruction of the extracellular matrix required for local and distal invasion. The proteolytic action of MMPs generates active matrix protein fragments, and influence the release, activation and bioavailability of growth factors. In addition, MMPs are involved in cell migration  
20   and in the processing or shedding of cell surface proteins. It is now believed that the proteolytic action of MMPs plays a very important role in tumor growth, apoptosis and angiogenesis (see Noel, A. et al., 'Emerging Roles for Proteinases in Cancer', Invasion Metastasis 17:221-239, 1997). Among the several MMPs known to date, gelatinase-A (MMP-2) is currently reputed as the main target in cancer.

25       Potent MMP inhibitors (MMPis) were disclosed three decades ago (e.g., see Dickens, J. P. et al., U.S. Patent 4,599,361), but only recently some of them entered clinical trials, and none has arrived at the market yet. Questions remain concerning their specificity, bioavailability, and potential long-term toxicity (see Hodgson, J.,

'Remodeling MMPs', Biotechnology 13:554-557, 1995), especially because they are primarily intended (in arthritis and cancer, at least) for prolonged therapy. Marimastat, the most advanced MMPi in the clinic, and other compounds currently investigated in patients such as Novartis CGS27023A and Agouron AG3340, have been reported to

5 cause a musculoskeletal syndrome characterized by joint pain and stiffness, which may be severe and dose-limiting. Although the cause of this toxicity is not well understood at this moment, it is thought to arise from an impairment in the normal tissue remodeling that is governed by one or more of the MMPs, primarily fibroblast collagenase (MMP-1). We have extended these observations by developing an animal model of joint

10 toxicity, involving treatment of rats for 10 days with MMPis at exposures close to that required for antitumor efficacy in mice (human prostate DU-145 model). In this rat model, MMPis characterized by poor selectivity (i.e., comparable potency against MMP-2 and MMP-1), such as marimastat and Roche Ro31-9790, elicited histological alteration of stifle joints (hypertrophic fibrosis of ligaments, interstitial hypertrophic


15 fibrosis of skeletal muscles, hypertrophic fibroplasia of the periosteum and synovium, chondrodysplasia, and decreased endochondral ossification of the epiphyseal plate); in some cases, clinical symptoms (abnormal posture, reluctance to move, abnormal gait) were also observed. Thus, there is a strong need for better MMPis, especially as far as the properties referred to above (specificity, bioavailability, long-term toxicity) are

20 concerned. The compounds of the present invention are intended to provide these advantages.

#### INFORMATION DISCLOSURE

A previous disclosure by the Pharmacia & Upjohn Company, PCT Patent Application WO 98/13340 (US Patent Number 5,847,153), provides a broad range of  $\beta$ -

25 sulfonyl hydroxamic acid derivatives, their preparation, and their activity against various enzymes of the MMP family, predominantly stromelysin and gelatinase. These compounds are generically described by the formula (Ia):




wherein  $R_1$  is  $C_{4-12}$  alkyl,  $C_{4-12}$  alkenyl,  $C_{4-12}$  alkynyl,  $-(CH_2)_h-C_{3-8}$  cycloalkyl,  $-(CH_2)_h$ -aryl, or  $-(CH_2)_h$ -het;  $R_2$  is  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $-(CH_2)_h-C_{3-8}$  cycloalkyl,  
 5  $-(CH_2)_h-C_{3-8}$  cycloalkenyl,  $-(CH_2)_h$ -aryl,  $-(CH_2)_h$ -het,  $-(CH_2)_h$ -Q,  $-(CH_2)_1$ -X- $R_4$ , or  $-(CH_2)_1$ - $CHR_5R_6$ .

We have now found that the combination of few particular groups  $R_1$  and  $R_2$  in a molecule of formula (Ia) provides distinct advantages over the other compounds of this class of MMPIs. Said combination, which is an object of the present invention, is  
 10 associated with outstanding potency ( $K_i$  in the picomolar range) against the gelatinases, in particular MMP-2, and relatively weak activity against fibroblast collagenase (MMP-1). Further, we have found that the same combination confers good potency (low nanoM range) and even higher selectivity for MMP-2 to the corresponding carboxylates, which were previously described as intermediates but not claimed as such. Finally, we have  
 15 found that the same combination provides advantages in the pharmacokinetic parameters, in particular lower clearance and higher oral bioavailability, and that antitumor efficacy in rodents can be achieved in the absence of joint toxicity.

#### DESCRIPTION OF THE INVENTION

The present invention provides 3-arylsulfonyl-2-methyl propanoic acid  
 20 derivatives of formula (I):



or pharmaceutically acceptable salts thereof wherein:

X is either HO-NH- (hydroxamic acids) or HO- (carboxylic acids);

5  $R_1$  is selected from phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-cyanophenyl, benzamido (i.e., -NH-CO-Ph) or benzamido substituted on the terminal Ph ring by  $C_{1-4}$  alkyl; fluoro, chloro, cyano,  $C_{1-4}$  alkoxy;

$R_2$  is selected from:

(a) -S-Ar or -S-CH<sub>2</sub>-Ar, wherein Ar is a monocarbocyclic or  
10 bicarbocyclic aromatic moiety, preferably phenyl, naphthyl and biphenyl, which is either unsubstituted or substituted with one or two substituents selected from the group consisting of  $C_{1-4}$  alkyl, phenyl, benzyl,  $C_{1-4}$  alkoxy, fluoro, chloro, bromo, nitro, cyano, hydroxy, amino, dimethylamino, acetamido, methylthio and acetyl;

(b) -O-Ar, wherein Ar is as defined above;


15 (c) -S-Het or -S-CH<sub>2</sub>-Het, wherein Het is a heterocyclic ring selected from the group consisting of pyridine, pyrimidine, pyridazine, pyrazine, 1,2,5-triazine, imidazole, thiophene, furan, pyrrole, pyrazole, 1,3-thiazole, 1,3-oxazole, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole, 1,2,3,4-tetrazole, quinoline, isoquinoline, indole, 1,3-benzoxazole, 1,3-benzothiazole, benzimidazole,  
20 [1,3]oxazolo[4,5-b]pyridine, [1,3]thiazolo[4,5-b]pyridine, [1,2,3,4]tetrazolo[1,5-b]pyridazine and purine, and wherein said Het group can be substituted with one to three substituents selected from the group consisting of  $C_{1-4}$  alkyl, phenyl, pyridyl, benzyl,  $C_{1-4}$  alkoxy, methylthio, fluoro, chloro, nitro, cyano, hydroxy, oxo, amino, methylamino, dimethylamino, 2-dimethylaminoethyl, acetamido and acetyl;

(d) 2,5-dioxo-1-imidazolidinyl or 2,4-dioxo-1-imidazolidinyl, which can be substituted at the carbon atom with one or two methyl groups, or with C<sub>2-4</sub> linear or branched alkyl, phenyl, benzyl or hydroxymethyl, and at the nitrogen atom with C<sub>1-4</sub> linear or branched alkyl.

5       The compounds of the present invention can be converted to pharmaceutically acceptable salts, where appropriate, according to conventional methods. The term 'pharmaceutically acceptable salts' refers to acid addition salts useful for administering the compounds of this invention and includes hydrochloride, sulfate, phosphate, acetate, propionate, lactate, mesylate, maleate, malate, succinate, tartrate, citrate, 2-hydroxyethyl  
10 sulfonate, fumarate and the like. Alternatively, the compounds of this invention, especially when X in formula (I) is hydroxy, may form metal salts, such as sodium, potassium, calcium or magnesium, or ammonium salts, or salts with an appropriate organic amine or amino acid such as arginine, procaine and the like. Any salt may be either in the anhydrous or hydrated form, or may be solvated with acceptable solvents,  
15 such as ethanol.

The present invention also provides pharmaceutical compositions which comprise, as active ingredient, a compound of formula (I) or a solvate, hydrate, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent or excipient.

20       The compounds of this invention contain a chiral center at the  $\alpha$ -position of the hydroxamic or carboxylic acid, and as such they may be obtained as any of the two separate enantiomers or as a racemic mixture of both. Preferred compounds of this invention are either the racemates or individual isomers having the configuration depicted in the formula (I') herebelow:



wherein X, R<sub>1</sub> and R<sub>2</sub> are as defined above. According to the Cahn-Ingold-Prelog rule, these preferred isomer are denoted as (*R*) when R<sub>2</sub> is -O-Ar, 2,5-dioxoimidazolidin-1-yl or 2,4-dioxoimidazolidin-1-yl, and as (*S*) when R<sub>2</sub> is -S-Ar, S-Het, -S-CH<sub>2</sub>-Ar or -S-CH<sub>2</sub>-Het. In addition, depending on the substituents, additional chiral centers or other isomeric forms may be present, and this invention embraces all possible stereoisomers and geometrical forms.

Particularly preferred compounds of this invention are as follows:

- 10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;  
 (2*S*)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;  
 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;  
 15 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[4-hydroxyphenyl)sulfanyl]methyl]propanoic acid;  
 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[[[4-hydroxyphenyl)sulfanyl]methyl]propanamide;  
 20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;  
 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;  
 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylsulfanyl)methyl]-propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(2-pyridinylmethyl)sulfanyl]methyl]-propanoic acid;

5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[[[(2-pyridinylmethyl)sulfanyl]-methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]-

10 methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[[[(5-methyl-1,3,4-thiadiazol-2-

yl)sulfanyl]methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(5-amino-1,3,4-thiadiazol-2-yl)-sulfanyl]-

15 methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(1-methyl-1H-imidazol-2-yl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfanyl)methyl]propanoic acid;

20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(1-methyl-1H-tetrazol-5-yl)sulfanyl]methyl]-

propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(1-benzyl-1H-tetrazol-5-yl)sulfanyl]methyl]-

25 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1H-benzimidazol-2-yl)sulfanyl)methyl]-propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1,3-thiazol-2-yl)sulfanyl)methyl]propanoic acid;

acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-methyl-2H-1,2,3-triazol-4-yl)sulfanyl]-

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[5-phenyl-1,3,4-oxadiazol-2-yl)sulfanyl]-

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[1-(3-pyridinyl)-1H-imidazol-2-yl)sulfanyl]-

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-methyl-1,3-oxazol-2-yl)sulfanyl)methyl]-

propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[1,3]oxazolo[4,5-b]pyrimidin-2-yl)sulfanyl]-

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[1,2,3,4]tetrazolo[1,5-b]pyridazin-6-yl)sulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2,6-dimethyl-4-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4,6-dimethyl-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-hydroxy-6-methyl-2-pyrimidinyl)sulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4,6-dihydroxy-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;



3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(9H-purin-6-ylsulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;

5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylaminophenoxy)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-acetylaminophenoxy)methyl]propanoic

10 acid;

3-[(1,1'-Biphenyl]-4-ylsulfonyl)-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]-propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid;

15 (2*R*)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-*N*-hydroxy-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-*N*-hydroxy-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

25 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2,5-dioxo-1-imidazolidinyl)-methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-methyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid;

5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-ethyl-2,5-dioxo-1-imidazolidinyl)methyl]-propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]-propanoic acid;

10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]-propanoic acid;

15 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-isopropyl-2,4-dioxo-1-imidazolidinyl)-methyl]propanoic acid;

20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-3-methyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;

25 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;  
3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfonyl)methyl]-propanamide;

3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;

- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-
- 5 propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
- 10 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylamino)phenoxymethyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;
- 15 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-pyridinylmethyl)sulfanyl]methyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 20 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 25 3-[[4-(Benzoylamino)phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]propanamide;
- 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[[2-pyridinylmethyl)sulfanyl]methyl]propanoic

acid;

3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid;

3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic  
5 acid;

3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-  
[(phenylsulfanyl)methyl]-

propanamide;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic  
10 acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-  
[(phenylsulfanyl)methyl]-

propanamide;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-  
15 propanamide;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic  
acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(2-  
20 pyridinylmethyl)sulfanyl)methyl]-

propanoic acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(1-methyl-1H-tetrazol-5-  
yl)sulfanyl]-

methyl]propanoic acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-  
25 imidazolidinyl)methyl]propanoic acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic  
acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-  
[(phenylsulfanyl)methyl]-

propanamide;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[[[(1-methyl-1H-tetrazol-5-  
5 yl)sulfanyl]-

methyl]propanoic acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-  
propanamide;

10 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic  
acid;

3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(3-  
fluorophenoxymethyl)-

propanamide;

15 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(2-acetyl amino)phenoxymethyl]-  
propanoic acid;


3-[[4-(4-Propylbenzoylamino)phenyl]sulfonyl]-2-[[[(1-methyl-1H-tetrazol-5-  
yl)sulfanyl]-

methyl]propanoic acid;

20 3-[[4-[(4-Methoxybenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(hydroxymethyl)phenoxy]methyl]-  
propanoic acid.

The compounds of this invention can be prepared by a process starting from a  
compound of formula 4:



wherein R is H or the residue of a carboxylic acid ester,  $R_1$  and  $R_2$  are as defined above and n is 0 or 2, which process comprises:

(a) hydrolysing a said compound of formula 4 in which R is the residue of a carboxylic and ester to give a compound of formula (I) in which X is HO-; or

(b) hydrolysing and oxidising, in either order, a said compound of formula 4 in which n is 0 and R is the residue of a carboxylic acid ester, to give a compound of formula (I) in which X is HO-; or


(c) activating a said compound of formula 4 wherein R is H and n is 2 to form an activated carboxy group; coupling the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotecting the hydroxamic group to give a compound of formula (I) wherein X is -NHOH; or

(d) submitting a said compound of formula 4 wherein R is H and n is zero to a sequence of reactions comprising oxidation at the sulphur atom, activation of the carboxy group, condensation of the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotection of the hydroxamic group to form a compound of formula (I) wherein X is -NHOH, the oxidation step being conducted either before the activation step or after the condensation step; and/or

(e) if desired, converting a resulting compound of formula (I) into another compound of formula (I); and/or converting a free compound into a pharmaceutically acceptable salt thereof; and/or converting a salt into a free compound.

The compound of formula 4 may be obtained by


(a) subjecting a compound of formula 2:




wherein R, R<sub>1</sub> and n are as defined above, to conjugate addition by treatment with a compound of formula R<sub>2</sub>H wherein R<sub>2</sub> is as defined above; or

(b) treating a compound of formula 3:

5



wherein R and R<sub>2</sub> are as defined above, with a thiol of formula:



10 to obtain a compound of formula 4 in which n is zero.

These processes are depicted in the following Scheme I, where R<sub>1</sub> and R<sub>2</sub> are the groups as defined previously, R is either a hydrogen atom or the residue of a carboxylic ester such as methyl, ethyl, n-butyl, tert-butyl, benzyl or benzhydryl, and n is either zero or 2. Compound 1, i.e. 2-(bromomethyl)acrylic acid or an ester thereof, is reacted with  
 15 the appropriate aryl-thiol in a suitable solvent such as toluene or dimethylformamide (DMF) in the absence or presence of a suitable base such as sodium bicarbonate or triethylamine (TEA), at ambient temperature or under heating up to reflux temperature, to afford a compound 2 wherein n is zero. The corresponding sulfone (i.e., a compound 2 wherein n is 2) is obtained, if desired, by subsequent reaction with an oxidant such as  
 20 metachloroperbenzoic acid (MCPBA) in a suitable solvent such as dichloromethane

(DCM) at 0 °C or ambient temperature, or using hydrogen peroxide in acidic acid; preferably, this reaction is performed with potassium peroxy monosulfate (Oxone<sup>®</sup>) in an organic solvent, preferably DMF, or in a mixture of water and water-miscible organic solvents such as methanol, DMF or the combination of two, at ambient temperature or under moderate heating. Alternatively, a compound 2 wherein n is 2 can be obtained from compound 1 directly, by reaction with the appropriate arylsulfinate. Preferably, the sulfinate is obtained and used as the sodium salt, or as the salt with a tertiary organic amine such as N-methylmorpholine (NMM). The sodium arylsulfinate can be conveniently obtained by reaction of the appropriate aryl-sulfonyl chloride with sodium iodide in acetone, preferably at ambient temperature, and it is isolated before reaction with compound 1, which takes place in a suitable solvent such as toluene, in the absence or presence of a suitable base such as sodium bicarbonate, at a temperature ranging from 25 °C to reflux. The salt with NMM is obtained by reaction of the arylsulfonyl chloride with 4-thiocresol and NMM in DCM, preferably at a temperature ranging from -50 to -78 °C, and it is reacted in situ with compound 1, then raising the temperature up to 25 °C or reflux.

A compound 4 can be obtained from the corresponding compound 2 by conjugate addition of a reagent of formula  $R_2H$ , or a salt thereof, where  $R_2$  is as defined previously. Thus, when  $R_2$  is a group  $-S-Ar$ ,  $-S-Het$ ,  $-S-CH_2-Ar$  or  $-S-CH_2-Het$ , the reagent is the corresponding thiol, or a salt thereof; when  $R_2$  is  $-O-Ar$ , the reagent is the corresponding aromatic carbinol; when  $R_2$  is 2,5-dioxo-1-imidazolidinyl or a substituted derivative thereof, the reagent is the corresponding 2,4-imidazolidinedione. Said conversion from a compound 2 into a compound 4 can be accomplished, under identical or similar conditions, either when n in 2 is zero or it is 2; however, the preparation of compounds of formula (I) wherein  $R_2$  is sulfanyl is better performed from compounds 2 where n is 2, because otherwise it is difficult to discriminate between the two sulfur atoms in the following oxidative step. Typically, the conjugate addition of said mercaptans and aromatic carbinols to a compound of formula 2 is accomplished using



the reagents neat, or in a suitable organic solvent such as DCM, DMF, toluene, tetrahydrofuran (THF), acetonitrile (MeCN), methanol or ethanol, at temperatures ranging from ambient to reflux, in the absence or presence of a suitable base such as sodium bicarbonate, sodium carbonate, TEA, sodium methoxide or sodium ethoxide, preferably under a nitrogen blanket. When the reagent is 2,4-imidazolidinedione or a substituted derivative thereof, the conjugate addition to a compound 2 is preferably performed in toluene or DMF in the presence of a suitable base such as sodium bicarbonate, TEA, NMM and the like, at a temperature ranging from ambient to reflux, preferably between 50 and 110 °C; alternatively, a preformed salt of said reagent can be used, especially sodium, potassium and lithium salts, which can be obtained by treatment with suitable strong bases, especially sodium hydride, potassium hexamethyldisilazide, potassium tert-butoxide and butyllithium, in a suitable organic solvent.

When a compound 4 where n is zero is obtained by the methods above, it may be converted, if desired, into the corresponding compound where n is 2 by oxidation, under the same conditions described previously for the oxidation of a compound 2 where n is zero to a compound where n is 2.

Alternatively, a compound 4 can be obtained from a compound 1 through the intermediacy of a compound 3. This alternative route provides advantages when the reagent  $R_2H$  is reluctant to the conjugate addition to a compound 2, e.g. when  $R_2H$  is 2,5-dioxo-1-imidazolidine, and particularly when it is 4,4-dimethyl-2,5-dioxo-1-imidazolidine and the N-alkyl derivatives thereof. In the first step, said reagent  $R_2H$  is condensed with the bromomethylacrylate 1 to provide 3. Typically, the reaction is performed in a solvent such as DMF or toluene, at temperatures ranging from 50 °C to reflux, and preferably in the presence of sodium bicarbonate. In the following step, conjugate addition of the suitable aryl-thiol to the intermediate 3 provides a compound 4 where n is zero. This reaction can be performed under the same conditions described above for reaction of 2 with a mercaptane  $R_2H$  to provide 4.

Compounds 4 where R is a hydrogen atom and n is 2 are specifically represented by structure 5 and as such are a first object of this invention, i.e. the compounds of formula (I) wherein X is hydroxy (the 'carboxylates'). Compounds 4 where R is the residue of a carboxylic ester, as defined previously, and n is 2, are converted to said  
 5 final compounds 5 by hydrolysis, according to the general methods known in the art such as using 6N HCl in acetic acid and refluxing for 10 to 20 hours or using iodotrimethylsilane in chloroform. When the R group is benzyl and R<sub>2</sub> is not a sulfanyl residue, a preferred method to obtain the free acid is hydrogenolysis, especially in the presence of a palladium catalyst such as Pd on charcoal, in an inert organic solvent such  
 10 as ethanol or DMF or the like, typically at room temperature and under atmospheric pressure or moderate pressure. When the R group is tert-butyl, a preferred method for obtaining the free acid is reaction with HCl or trifluoroacetic acid, in inert organic solvents such as DCM, THF, MeCN and the like, optionally in admixture with water, methanol or ethanol, at temperatures ranging from 20 to 40 °C.

15 Compounds 4 where R is the residue of a carboxylic ester and n is zero are converted to compounds 5 by a two-step procedure involving oxidation and hydrolysis, or hydrolysis and oxidation, as preferred. Conditions for oxidation and hydrolysis are the same as described above.

Compounds 4 where R is a hydrogen atom and n is 2, which have structure 5,  
 20 are converted to compounds 6, which specifically represent the second object of this invention, i.e. structure (I) wherein X is -NHOH (the 'hydroxamates'), by activation of the acid and coupling with hydroxylamine or an O-protected derivative thereof. This reaction can be performed by several routes well known to those skilled in the art. For example, the acid can be activated as its chloride or a mixed anhydride. A preferred  
 25 method for activation as the acid chloride is reaction with oxalyl chloride, typically in DCM containing DMF as a catalyst, at 0 °C to ambient temperature. A preferred method for activation as a mixed anhydride is reaction with a chlorocarbonate such as chloroethylformate in anhydrous THF or a similar compatible solvent, in the presence of


a tertiary amine such as TEA, NMM and the like. Alternatively, the acid can be activated by a carbodiimide condensing agent such as dicyclohexylcarbodiimide (DCC) or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (WSCDI, water soluble carbodiimide), preferably in the presence of 1-hydroxybenzotriazole (HOBt) in an inert organic solvent such as DMF at 0 °C to room temperature. Another method for the activation of the acids of formula 4 where R is hydrogen is reaction with benzotriazolyl-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent) in a suitable organic solvent such as MeCN, THF or DCM in the presence of a tertiary amine such as TEA or NMM. A further method is reaction with O-benzotriazolyl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) or O-benzotriazolyl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), under conditions similar to those described for the BOP reagent. When the activation is performed with oxalyl chloride, the acid chloride is usually isolated crude before reaction with the hydroxylamine reagent; when with chloroethylformate, EDC or WSCDI, the hydroxylamine reagent is preferably added after about 1 hour at room temperature; when with the BOP reagent, HBTU or TBTU, the hydroxylamine reagent is O-protected and it is preferably added from the beginning. Suitable O-protected hydroxylamines are O-(benzyl)benzylhydroxylamine, O-(tert-butyl)hydroxylamine, O-(tetrahydropyranyl)-hydroxylamine, O-(trimethylsilyl)hydroxylamine and O-(tert-butyl)dimethylsilyl)-hydroxylamine, the first two being preferably used as their hydrochlorides and freed in situ by a tertiary amine such as TEA or NMM. Alternatively, when activation of the acid is performed with oxalyl chloride or WSCDI, the unprotected hydroxylamine (e.g., as the commercial 50% solution in water) or the hydrochloride thereof can be used, either in the organic solvents specified above or after dissolution in water or in a mixture of water and THF. Suitable conditions for the condensation step vary according to the activation method. Typically, when activation is by the acid chloride method, the reaction is fast and is complete within 1-5 hours at temperatures ranging from 0 °C to 25 °C; otherwise it can be slower, usually requiring

stirring overnight, at ambient temperature or even 1-5 days. When an O-protected hydroxylamine is used in the condensation, the reaction affords O-protected hydroxamate derivatives, and the desired free-hydroxamates 6 are obtained after removal of the protecting group. In particular, when O-(benzyl)hydroxylamine is used, deprotection is performed by hydrogenolysis, preferably with a palladium catalyst such as 5% Pd on charcoal, 5% Pd on BaSO<sub>4</sub>, or palladium hydroxide on carbon (Pearlman's catalyst), typically in methanol or ethanol. When O-(tert-butyl)hydroxylamine is used, the free hydroxamate can be obtained after treatment with neat or aqueous trifluoroacetic acid, or with HCl in DCM. When O-(tetrahydropyranyl)hydroxylamine is used, the free hydroxamate is rapidly obtained by reaction with 4N aqueous HCl in suitable organic solvents or mixtures thereof, e.g. dioxane and methanol. When O-(trimethylsilyl)hydroxylamine is used, deprotection is usually achieved during work up, e.g. by partitioning of the condensation mixture between ethyl acetate and diluted hydrochloric acid. When O-(tert-butyl dimethylsilyl)hydroxylamine is used, deprotection can be achieved by HF or fluoride salts, preferably with tetrabutylammonium fluoride, in a suitable organic solvent such as THF, at temperatures ranging from -10 to +25 °C.

Compounds 4 where R is a hydrogen atom and n is zero are converted to compounds 6 by a sequence involving oxidation at the sulfur atom, activation of the carboxy group, condensation of the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, when required, unmasking of the free hydroxamic group. The oxidation can be performed first, i.e. on said compounds of formula 4, as described previously for the oxidation of a compound of formula 2 where n is zero to the corresponding compound where n is 2. Alternatively, the oxidation can be performed at a later stage, i.e. on the O-protected or free hydroxamic acid, in which case a preferred oxidant is oxone, in a suitable organic solvent such as DMF, mixture of water and one or two water miscible organic solvents, such as methanol, ethanol, DMF and THF, at a temperature ranging from 5 to 25 °C. Conditions for the other steps of the sequence, i.e. activation of the carboxy group, coupling with hydroxylamine or a

protected derivative thereof and, if required, deprotection, are the same as described previously.

# SCHEME I



In some cases, minor modifications of the general procedure indicated in Scheme 1 can be used to obtain compounds 4. For example, conversion of a compound 2 into a compound 4 wherein  $\text{R}_2$  is a sulfanyl residue can be achieved in two steps, the first involving addition of HBr to a compound 2 to obtain the bromomethyl derivative, which is then reacted with the thiol which  $\text{R}_2\text{H}$  can represent. This two-step procedure


10

can be convenient for less reactive thiols, in particular when  $R_2$  is  $-S-CH_2-Ar$  or  $-S-CH_2-Het$ . A preferred solvent for the addition of  $HBr$  to a compound 2 is acetic acid, usually at ambient temperature. The thiol is preferably reacted with said bromomethyl intermediate in the presence on an organic or inorganic base such as TEA, NMM, sodium bicarbonate, potassium carbonate and the like. Less reactive thiols can be advantageously reacted after conversion to salts thereof, e.g. by treatment with potassium hexamethyldisilazide at low temperature in a suitable organic solvent such as THF, and reacted in situ with the compound 2. The bromomethyl intermediate, or the corresponding carbinol, can also be used to obtain compounds 4 wherein  $R_2$  is 2,5-dioxo-1-imidazolidinyl. In particular, the bromomethyl derivative can be reacted with 2,4-imidazolidinedione, or a derivative thereof, under conditions similar to that described above for reaction with a thiol, and the carbinol can be condensed with the same reagent under Mitsunobu conditions, i.e. in the presence of triphenylphosphine and diethyl azodicarboxylate, in DMF at temperatures ranging from 0 to 25°C.

The compounds of formula (I) wherein  $R_2$  is 2,4-dioxo-1-imidazolidinyl can be obtained by the variant described in Scheme II below. As therein outlined, an amino acid  $H_2N-CHR_3-COOH$  where  $R_3$  is a hydrogen atom,  $C_{1-4}$  straight or branched alkyl or benzyl, or a salt or an ester thereof, especially the methyl or ethyl esters, is condensed with compound 1, as defined in Scheme I, to obtain a compound 7 where  $R$ ,  $R_3$  are as above defined, and  $R'$  is either a hydrogen atom or methyl or ethyl. This reaction is carried out in a suitable organic solvent, with or without a base, at temperatures ranging from 0 to 50 °C, preferably around ambient temperature. A preferred solvent is DMF; preferred bases, which can be present in amounts variable to a catalytic one to 3 molar equivalents, are tertiary amines, such as TEA, NMM, Hunig's base and the like. The obtained compound 7 is then reacted with sodium or potassium cyanate, or with an isocyanate of formula  $R_4-NCO$  where  $R_4$  is  $C_{1-4}$  alkyl, phenyl or benzyl, to obtain an intermediate urea, which is cyclized to a compound 8 where  $R$ ,  $R_3$  and  $R_4$  are as defined above, in a suitable organic solvent such as THF, dioxane, MeCN, DMF and the like, at

temperatures ranging from about 20 °C to reflux. Said compound 8 is then reacted with an aryl thiol of formula  $R_1-C_6H_4-SH$  to obtain a compound 9 where  $R_1$  is as above defined and  $n$  is zero. Typically, the conjugate addition of said aryl thiol to a compound 8 is accomplished in a suitable organic solvent such as DCM, DMF, toluene, tetrahydrofuran (THF), acetonitrile (MeCN), methanol or ethanol, at temperatures ranging from ambient to reflux, in the absence or presence of a suitable base such as sodium bicarbonate, sodium carbonate, TEA, sodium methoxide or sodium ethoxide, preferably under a nitrogen blanket. Conversion of a compound 9 where  $n$  is zero into a compound 9 where  $n$  is 2 is accomplished under the conditions previously described in Scheme I, preferably with Oxone® in an organic solvent such as DMF or in a mixture of water and water-miscible organic solvent such as methanol, DMF or the combination of two, at ambient temperature or under moderate heating. Alternatively, the same synthetic sequence is carried out on a compound 2, as defined in Scheme I, to obtain the same compound 9 *via* 10. Compounds 9 where  $n$  is two and  $R$  is a hydrogen atom are the desired compounds of formula (I) where  $X$  is HO- and  $R_2$  is 2,4-dioxoimidazolidin-1-yl. The corresponding compounds of formula (I) where  $X$  is HONH- are obtained, if desired, as described previously (Scheme I) for the conversion of a compound 5 into a compound 6.

## SCHEME II



When required or convenient, chemical modifications of the group  $R_1$  or  $R_2$  in intermediates 2, 3, 4 and 9 can be performed in the above processes to obtain the final compounds of formula (I). For example, a compound 3 or 4 wherein  $R_2$  is 2,5-dioxo-1-imidazolidinyl having the 3-nitrogen atom unsubstituted can be alkylated at said atom to obtain the corresponding N-alkyl derivatives, e.g. with methyl iodide in DMF in the presence of potassium carbonate. When  $R_1$  and  $R_2$  in the final compounds of formula (I) are, or contain, a reactive functional group, e.g. amino and hydroxy, these groups can be protected at the appropriate stage and and unmasked when convenient, according to conventional methods well known in the art.



The chemistry in Scheme I and II proceeds through achiral or racemic intermediates, unless a substituent in  $R_1$  or  $R_2$  is chiral and enantiopure, or the aminoacid used in Scheme II for the preparation of compound 7 or 10 is chiral and enantiopure. When desired, pure enantiomers of the final products, in particular compounds of formula (I') above, may be obtained from racemic compounds (I), or racemic intermediates thereof, by methods well known to one skilled in the art. For example, racemic carboxylates, including compounds (I) where X is HO- and intermediates 4, 9 where R is a hydrogen atom and n is zero, or intermediates 10 where R is a hydrogen atom, can be resolved by chiral salt formation, for example with enantiopure 1-phenylethylamine, brucine, ephedrine, strychnine, morphine. Alternatively, final racemic compounds or racemic intermediates can be resolved by chiral chromatography. By another method, carboxylic esters, preferably methyl, ethyl and butyl esters, can be selectively hydrolysed to chiral acids by lipases and esterases. By a still different procedure, racemic carboxylic acids can be converted to diastereomeric esters with chiral alcohols such as borneol, menthol and the like, which are then separated by crystallization or conventional chromatography. Alternatively, when in the acrylates of formula 2, 3, 8 in Schemes I and II R is the residue of a chiral alcohol such as borneol, menthol and the like, diastereomeric compounds 4, 9 10 are obtained, and variable degrees of diastereoselectivity can be achieved.

The present invention further provides a compound of formula (I) as defined above for use in a method of treatment or prophylaxis of a disease mediated in a mammal by a matrix metalloproteinase, especially the gelatinases (MMP-2 and MMP-2), the membrane-type MMP involved in gelatinase activation (MMP-14), the stromelysins (MMP-3 and MMP-10), collagenase-3 (MMP-13) and neutrophyl collagenase (MMP-8). Diseases for which the compounds of the present invention are particularly intended are: tumor growth and metastasis; rheumatoid arthritis and osteoarthritis; ophthalmic diseases, in particular age-related macular degeneration and

diabetic retinopathy; cardiovascular diseases, in particular, restenosis; periodontal diseases; multiple sclerosis; Alzheimer disease.

The compounds of the present invention are typically administered in the form of pharmaceutical compositions, which can be prepared by combining the compounds of formula (I) or (I') or salts or hydrates or solvates thereof with a solid or liquid pharmaceutically acceptable carrier, and, optionally, with pharmaceutically acceptable adjuvants and excipients, employing standard and conventional techniques. In particular, said compounds can be administered:

a) orally, for example, as tablets, coated tablets, capsules, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is present as such, or mixed with water or an oil medium, for

example, peanut oil, liquid paraffin, or olive oil. Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, 5 polyvinylpyrrolidone gum tragacanth and gum acacia; dispersing or wetting agents may be naturally-occurring phosphatides, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial 10 esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The said aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, or one or 15 more sweetening agents, such as sucrose or saccharin. Oily suspension may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to 20 provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid. Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are 25 exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oils, or a mineral oil, for

example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan mono-oleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, for example glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents;

b) parenterally, either subcutaneously, or intravenously, or intramuscularly, or intrasternally, or by infusion techniques, in the form of sterile injectable aqueous or oleaginous suspensions. This suspension may be formulated according to the known art using those suitable dispersing, wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition fatty acids such as oleic acid find use in the preparation of injectables;

c) topically, in the form of creams, ointments, jellies, plasters, collyriums, solutions or suspensions;


d) rectally, in the form of suppositories prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperature but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

The compounds of this invention are advantageously administered orally in solid and liquid dosage forms. Daily dosages can vary within wide limits and will be adjusted to the individual requirements in each particular case. As a rule of thumb, a compound of formula (I) or (I') is advantageously administered at a dosage and frequency to achieve and maintain, for the whole period of therapy, a blood level in free drug (i.e., not bound to plasma proteins) ranging from about 1 to about 100 times its  $K_i$  against the most relevant MMP for the particular disease state, e.g. MMP-2 for cancer. Thus, the daily dose may be divided into multiple doses for administration, e.g. two to four times per day, which is generally expedient for compounds of formula (I) or (I') wherein X is HO-NH-; or, in the case of particularly favorable pharmacokinetics, which can be found in compounds of formula (I) or (I') wherein X is HO-, the drug can be given daily, or even at alternate days. Generally, an amount of the active compound effective in adult patients will be in the range of 1 to about 500 mg, but it is to be understood that such range may be exceeded depending on the particular compound being used, the severity of the disease being treated, and the requirement of the patient.

The compounds of the present invention and their preparation will be better understood in connection with the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

#### EXAMPLE 1

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid.



Step 1. 4'-Chloro[1,1'-biphenyl]-4-sulfonic acid.

To a stirred solution of 4-chlorobiphenyl (21 g) in 150 mL of chloroform, cooled at  $-5^{\circ}\text{C}$ , was added dropwise chlorosulfonic acid (10.2 mL, 1.36 eq). The cooling bath was removed and the mixture was stirred for 1 hour. Evaporation in vacuo left the title compound as a syrup.

5 Step 2. 4'-Chloro[1,1'-biphenyl]-4-sulfonyl chloride.

The crude product from step 1 was dissolved in thionyl chloride (100 mL) and refluxed for 1 hour. The solvent was removed in vacuo and the resulting solid was stripped twice with toluene (150 mL).

Step 3. 4'-Chloro[1,1'-biphenyl]-4-thiol.

10 4'-Chloro[1,1'-biphenyl]-4-sulfonyl chloride (30 g) was added portion wise to a stirred solution of tin(II) chloride dihydrate (80 g) in conc. HCl (50 mL) and water (150 mL) at  $90-95^{\circ}\text{C}$ . The resulting yellow slurry was stirred for 1 hour at the same temperature. After cooling, the precipitate was collected by filtration, thoroughly washed with water and dried to afford 21 g of bis[4-(4'-chloro[1,1'-biphenyl])]disulfide, mp  $164-166^{\circ}\text{C}$ .  
 15 This material was dissolved in THF (200 mL) and conc. HCl (25 mL), and Zn powder (20 g) was added under stirring. After stirring for 30 minutes at ambient temperature, the Zn was removed and rinsed with THF. The clear solution was concentrated in vacuo and partitioned between chloroform (450 mL) and brine. The organic phase was dried over anhydrous sodium sulfate and evaporated to leave 19 g of the title compound, mp  
 20  $154-157^{\circ}\text{C}$ .

Step 4. 2-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]acrylic acid.

The thiol from step 3 (22 g) and 2-(bromomethyl)acrylic acid (18 g) were dissolved in 100 mL of DMF and stirred at  $30^{\circ}\text{C}$  for 2 days under a low stream of nitrogen. The reaction mixture was diluted with ethyl acetate (400 mL) and washed  
 25 several times with brine until neutral. The organic phase was dried over anhydrous sodium sulfate and evaporated in vacuo to obtain a crystalline mass. After addition of diethyl ether (150 mL) and swirling, the solid was collected by filtration, washed twice with ether, and dried to afford 21 g of the title compound, mp  $191-195^{\circ}\text{C}$ .

Step 5. 2-[[4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]acrylic acid.

The material from step 4 above (20 g) was dissolved in 100 mL of DMF and added dropwise to a stirred solution of Oxone® (105 g) in 300 mL of water at 45-50 °C. After stirring for 5 hours, water (1 L) was added. The solid was collected by filtration,  
 5 washed several times with water, then twice with ethanol (100 mL) and dried to provide 21.5 g of the title compound, mp 178-181 °C.

Step 6. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid.

Under a low stream of nitrogen, thiophenol (20 mL) was added to a stirred  
 10 solution of the material from step 5 above (20 mg) in 35 mL of DMF. After stirring at 35 °C for 2 days, the reaction mixture was diluted with ethyl acetate and treated dropwise with a solution of iodine (21 g) in 75 mL of THF. The resulting brown solution was washed until complete decolorization with a solution of sodium metabisulfite (10 g) in 300 mL of water. After washing with brine and drying over  
 15 anhydrous sodium sulfate, the solvent was removed and the residue was filtered through a short pad of silica gel (450 g), eluting with chloroform, to obtain 23 g of the title compound, mp 158-162 °C.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 3.06 (m, 2H), 3.33 (m, 1H), 3.60 (dd, *J*= 3.4 and 14.2 Hz, 1H), 3.68 (dd, *J*= 7.3 and 14.2 Hz, 1H), 7.22 (m, 5H), 7.50 (m, 4H), 7.68 (d, *J*=  
 20 8.4 Hz, 2H), 7.88 (d, *J*= 8.4 Hz, 2H) [COOH: not seen]. MS (EI): *m/z* 446, 252, 194, 152, 149, 110. Anal. Found C, 59.32; H, 4.44; S, 14.32.

#### EXAMPLE 2

Preparation of sodium 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoate.

25 Sodium hydroxide (1 M, 48 mL) was added to a solution in methanol (75 mL) of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2(*RS*)-[(phenylsulfanyl)-methyl]propanoic acid (20 g, obtained as described in Example 1). Charcoal was added and the mixture was filtered. Concentration of the filtrate to 1/3 of the original volume furnished a shiny

precipitate, which was collected by filtration and sequentially washed with cold water, ethanol and diethyl ether. Drying to a constant weight provided 1.9 g of the title compound as white crystals.

### EXAMPLE 3

5 Preparation of (2*S*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid and of (2*R*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid.

Racemic 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid, prepared as described in Example 1, was eluted over Chiralpak AS  
10 (Chiral Technologies) with ethanol + 0.05% trifluoroacetic acid (V/V) at 0.4 mL/min, monitoring the absorbance at 270 nm. The eluates were collected and concentrated in vacuo to yield the title enantiomers.

ISOMER 1, (2*S*) configuration:


$R_f = 12.0$  min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA);  $[\alpha]_D^{25} =$   
15  $+47.8^\circ$  ( $c = 0.83\%$ , EtOH). Absolute configuration established by X-Ray crystallography.

ISOMER 2, (2*R*) configuration:

$R_f = 15.0$  min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA);  $[\alpha]_D^{25} = -43^\circ$   
( $c = 0.83\%$ , EtOH).

### EXAMPLE 4

20 Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfonyl)methyl]propanamide.






TBTU (0,72 g) was added portionwise to a stirred solution of N-methylmorpholine (0.5 mL) and O-tert-butylhydroxylamine hydrochloride (0.43 g) and 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid (0.5 g) in 20 mL of DMF and 20 mL of acetonitrile cooled at -5°C. After stirring at ambient temperature overnight, the resulting solution was diluted with ethyl acetate and sequentially washed with 0.1 M citric acid and 0.1 M NaHCO<sub>3</sub>. After drying over anhydrous sodium sulfate and removal of the solvent, the crude reaction mixture was treated with trifluoroacetic acid (10 mL) and kept aside for 6 hours. The solvent was removed and the residue was partitioned between ethyl acetate and 0.1 M NaHCO<sub>3</sub>. After drying, the solvent was evaporated off and the residue was chromatographed on silica gel eluting with chloroform/cyclohexane 2/3 to afford, after crystallization from ethyl acetate, 0.35 g of the title compound, mp .....

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.70 (m, 1H), 3.09 (d, *J*= 7.8 Hz, 2H), 3.61 (m, 2H), 7.20 (m, 5H), 7.58 (d, *J*= 8.3 Hz, 2H), 7.78 (d, *J*= 8.3 Hz, 2H), 7.90 (m, 4H), 8.95 (s, 1H), 10.81 (s, 1H).. MS (EI): *m/z* 461, 400.

#### EXAMPLE 5

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.



Step 1. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.

To a stirred solution of α-bromomethylacrylic acid (1.65 g) and NaHCO<sub>3</sub> (1.7 g) in 25 mL of DMF was added 3,4,4-trimethyl-2,5-dioxo-imidazolidine (2.13 g) and the

resulting suspension was heated at 75°C for 3 h. After disappearance of  $\alpha$ -bromomethylacrylic acid, checked by tlc and HPLC, the solution was cooled at 0-5°C and 4'-chloro[1,1'-biphenyl]-4-thiol (3.3 g) was added. After stirring overnight at room temperature, the solution was diluted with ethyl acetate and washed thoroughly with 0.1 M citric acid. After drying and removal of the solvent, the residue was chromatographed on silica gel eluting with chloroform/cyclohexane 1/1 to provide, after crystallization from ethyl ether, 2.4 g of the title compound, mp 148-150 °C.

Step 2. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.

A solution of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid (4.6 g) in 30 ml of DMF was added dropwise to a stirred solution of Oxone® (13.5 g) in 75 ml of DMF and 20 mL of water at 45°C. After stirring for 2 h, the solvent was removed and the residue was partitioned between ethyl acetate and brine. The residue was columned on a small pad of silica gel eluting with chloroform to furnish 3.9 g of the title compound as a white foam.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  1.22 (s, 3H), 1.24 (s, 3H), 2.74 (s, 3H), 3.02 (m, 1H), 3.5-3.7 (m, 4H), 7.57 (d, *J* = 8.3 Hz, 2H), 7.79 (d, *J* = 8.3 Hz, 2H), 7.94 (m, 4H), 12.85 (broad s, 1H). MS (EI): *m/z* 478, 433, 252, 211, 152, 56. Anal. Found C, 54.89; H, 5.15; N, 5.64; S, 6.53.

#### EXAMPLE 6

Preparation of (2*R*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid and of (2*S*)-3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.

Racemic 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid, prepared as described in Example 5, was separated by chiral chromatography under the conditions described in Example 3.

ISOMER 1, (2*R*) configuration:


$R_f = 9$  min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA);  $[\alpha]^{25}_D = -8.7^\circ$  ( $c = 0.83\%$ , EtOH). The absolute configuration established by protein X-Ray in complex with truncated stromelysin (MMP-3).

ISOMER 2, (2*S*) configuration:

- 5  $R_f = 12$  min (0.46x25 cm Chiralpak AS, ethanol + 0.05% TFA);  $[\alpha]^{25}_D = +8.2^\circ$  ( $c = 0.83\%$ , EtOH).

#### EXAMPLE 7

Preparation of 3-([1,1'-biphenyl]-4-ylsulfonyl)-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.




Following the general procedure outlined in Example 5 and making non-critical variations but replacing 4'-chloro[1,1'-biphenyl]-4-thiol with [1,1'-biphenyl]-4-thiol in step 2, the title compound was obtained as a whitish crystalline powder, mp > 200 °C (dec.).

$^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  1.16 (s, 3H), 1.17 (s, 3H), 2.68 (s, 3H), 2.96 (m, 1H), 3.43 (dd,  $J = 3.6$  and 14.6 Hz, 1H), 3.54 (m, 2H), 3.63 (dd,  $J = 8.7$  and 14.6 Hz, 1H), 7.40 (m, 3H), 7.70 (m, 2H), 7.87 (m, 4H), 12.90 (broad s, 1H). MS (EI):  $m/z$  445, 427, 227, 209. Anal. Found C, 57.39; H, 5.66; N, 6.98; S, 5.95.

#### EXAMPLE 8

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide.




Following the general procedure outlined in Example 4 and making non-critical variations but replacing 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid with 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid, the title compound was obtained as a white crystalline powder.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.19 (s, 3H), 1.22 (s, 3H), 2.72 (s, 3H), 2.90 (m, 1H), 3.38 (dd, *J* = 2.9 and 14.6 Hz, 1H), 3.46 (m, 2H), 3.60 (dd, *J* = 8.3 and 14.6 Hz, 1H), 7.57 (d, *J* = 8.3 Hz, 2H), 7.78 (d, *J* = 8.3 Hz, 2H), 7.93 (m, 4H), 8.82 (s, 1H), 10.80 (s, 1H). ESI(+) MS: *m/z* 494, 461, 235, 209. Anal. Found: C, 43.58; H, 4.42; N, 6.61; S, 5.10.

#### EXAMPLE 9

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.




Following the general procedure outlined in Example 5 and making non-critical variations but starting with 5,5-dimethyl-2,4-imidazolidinedione in step 1, the title compound was obtained as a white crystalline powder, mp 242-245 °C.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO-}d_6$ ):  $\delta$  1.21 (s, 3H), 1.22 (s, 3H), 3.04 (m, 1H), 3.52 (m, 1H), 3.57 (m, 1H), 3.68 (m, 1H), 7.58 (d,  $J = 8.5$  Hz, 2H), 7.79 (d,  $J = 8.5$  Hz, 2H), 7.94 (m, 4H), 8.29 (s, 1H), 12.82 (broad s, 1H). ESI(+)-MS:  $m/z$  465, 235, 213, 213, 195. Anal. Found C, 52.30; H, 4.82; N, 5.76; S, 6.44.

5

## EXAMPLE 10

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide.



10

Following the general procedure outlined in Example 7 and making non-critical variations but starting in step 1 with 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid (prepared as described in Example 8), the title compound was obtained as a white crystalline powder.


15

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO-}d_6$ ):  $\delta$  1.17 (s, 3H), 1.21 (s, 3H), 3.34 (dd,  $J = 3.0$  and 15.0 Hz, 1H), 3.44 (m, 2H), 3.62 (dd,  $J = 9.0$  and 15.0 Hz, 1H), 7.58 (d,  $J = 8.9$  Hz, 2H), 7.78 (d,  $J = 8.9$  Hz, 2H), 7.93 (m, 4H), 8.24 (s, 1H), 8.83 (s, 1H), 10.81 (s, 1H). ESI(+)-MS:  $m/z$  480, 447, 195. Anal. Found C, 54.06; H, 4.87; N, 8.67; S, 6.96.

## EXAMPLE 11

20

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid.




Following the general procedure outlined in Example 5 and making non-critical variations but starting with 2,4-imidazolidinedione in step 1, the title compound was obtained as a white crystalline powder, mp 188-190 °C.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 3.02 (m, 1H), 3.5-3.7 (m, 4H), 3.82 (d, *J*= 17.7 Hz, 1H), 3.87 (d, *J*= 17.7 Hz, 1H), 7.59 (d, *J*= 8.5 Hz, 2H), 7.80 (d, *J*= 8.5 Hz, 2H), 7.94 (m, 4H), 8.07 (s, 1H), 12.80 (broad s, 1H). ESI(+) MS *m/z* 437, 419, 235, 167. Anal. Found: C, 47.71%; H, 3.81%; N, 5.79%; S, 7.18%.

#### EXAMPLE 12

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanamide.




Following the general procedure outlined in Example 8 and making non-critical variations but starting in step 1 with 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid (prepared as described in Example 11), the title compound was obtained as a yellowish crystalline powder, mp 150-156 °C.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.83 (m, 1H), 3.37 and 3.60 (each m, 1H), 3.42 (m, 2H), 3.76 (d, *J*= 17.6 Hz, 1H), 3.81 (d, *J*= 17.6 Hz, 1H), 7.57 (d, *J*= 8.3 Hz,

2H), 7.78 (d,  $J$ = 8.3 Hz, 2H), 7.91 (m, 4H), 8.01 (s, 1H), 8.70 (broad s, 1H), 10.80 (broad s, 1H). ESI(+)-MS:  $m/z$  452, 419, 235. Anal. Found C, 44.35; H, 3.87; N, 7.59; S, 6.81.

## EXAMPLE 13

- 5 Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid.



Step 1. Methyl 2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]acrylate.

Methyl 2-(bromomethyl)acrylate (0.7 mL) and 1-butyl-2,4-imidazolidinedione (1 g) were dissolved in DMF (3 mL). Sodium hydrogen carbonate (698 mg) was added, and the mixture heated at 80 °C for 2 hours, cooled to ambient temperature and partitioned between ethyl acetate and water. The organic layer was washed with water  
 5 and brine, dried over anhydrous sodium sulfate and evaporated in vacuo to leave the title compound (1.0 g) as a colourless oil.

Step 2. Methyl 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoate.

The product from step 1 above (500 mg) and 4'-chloro[1,1'-biphenyl]-4-thiol  
 10 (520 mg) were dissolved in DMF (5 mL). Sodium hydrogen carbonate (252 mg) was added, and the mixture heated at 60 °C for 1 hour, cooled to ambient temperature and partitioned between ethyl acetate and water. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and evaporated in vacuo to leave the crude title compound, which was used as such in the following step.

15 Step 3. Methyl 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoate.

The crude compound from step 2 above was taken up in DMF (7 mL). Oxone® (2 g) was added, and the mixture stirred for 4 hours at ambient temperature. Work-up as in step 1 left a residue, which was purified by flash-chromatography over silica (ethyl  
 20 acetate – hexane), to obtain the title compound (650 mg) as a colorless foam.

Step 4. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)-methyl]propanoic acid.

2N Hydrochloric acid (5 mL) was added to a solution of the compound from step 3 above (650 mg) in acetic acid (2 mL), and the mixture was stirred at 80 – 90 °C.  
 25 After 2 hours, 37% HCl (1 mL) was added and stirring at 80 °C was continued for additional 5 hours. The reaction mixture was evaporated in vacuo, and the residue partitioned between ethyl acetate and water. The organic layer was dried over anhydrous sodium sulfate and evaporated. The solvent was removed in vacuo and the




resulting solid was stripped twice with toluene, to yield the title compound (420 mg) as a whitish powder.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 0.84 (t, *J* = 7.3 Hz, 3H), 1.07 (m, 2H), 1.41 (m, 2H), 2.99 (m, 1H), 3.20 (t, *J* = 7.1 Hz, 2H), 3.4-3.7 (m, 4H), 3.88 (d, *J* = 7.6 Hz, 1H),  
 5 3.92 (d, *J* = 7.6 Hz, 1H), 7.58 (d, *J* = 8.8 Hz, 2H), 7.80 (d, *J* = 8.8 Hz, 2H), 7.93 (m, 4H),  
 12.95 (broad s, 1H). ESI(+)-MS: *m/z* 515, 493, 475, 223. Anal. Found C, 57.39; H, 5.66; N, 6.98; S, 5.95.

#### EXAMPLE 14

Preparation of 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid.



Step 1. Bis(4-(benzoylamino)phenyl) disulfide.

Benzoyl chloride (8 mL) was added to a solution of bis(4-aminophenyl) disulfide  
 15 (4 g) and TEA (10 mL) in 100 mL of dichloromethane, and the mixture was stirred for 2 hours at ambient temperature. The precipitate was collected by filtration and sequentially washed with water, diluted aqueous HCl and n-hexane to obtain the title compound as a white solid.

Step 2. N-(4-Sulfanylphenyl)benzamide.

20 Zn powder (0.5 g) and 2 N HCl (30 mL) was added in three portions under vigorous stirring to a suspension of the disulfide from step 1 in THF (100 mL). After 4 hour at ambient temperature, the mixture was filtered and the filtrate was concentrated in vacuo. The residue was taken up in ethyl acetate and sequentially washed with water and

aqueous sodium hydrogen carbonate. Drying over anhydrous sodium sulfate and evaporation afforded 3.66 g of the title compound as a yellow solid.

Step 3. 2-[[[4-(Benzoylamino)phenyl]sulfonyl]methyl]acrylic acid.

A mixture of N-(4-sulfanylphenyl)benzamide (1.5 g), 2-(bromomethyl)acrylic acid (1 g) and TEA (1.75 mL) was stirred for 4 hours at ambient temperature. Washing with diluted aqueous HCl, drying over anhydrous sodium sulfate and evaporation in vacuo left a residue, which was passed through a short pad of silica gel (dichloromethane – methanol as the eluants) to obtain 1.68 g of the title compound as a colorless oil.

Step 4. 2-[[[4-(Benzoylamino)phenyl]sulfonyl]methyl]acrylic acid.

A mixture of Oxone<sup>®</sup> (10 g) and 2-[[[4-(benzoylamino)phenyl]sulfonyl]methyl]acrylic acid (1.68 g) in DMF (20 mL) was heated at 50 °C for 4 hours under vigorous stirring, then partitioned between ethyl acetate and 4% aqueous HCl. The organic phase was washed twice with water, dried over anhydrous sodium sulfate and evaporated to obtain the crude title compound as a yellowish oil.


Step 5. 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid.

2-Thiophenethiol (250 mg) was added to a solution of the crude acrylic acid from step 4 above (300 mg) in 5 mL of DMF. After 1 hour at ambient temperature, the mixture was partitioned between ethyl acetate and 4% aqueous HCl. The organic phase was washed twice with water, dried over anhydrous sodium sulfate and evaporated. The residue was chromatographed over silica gel with chloroform/methanol, and the eluate concentrated in vacuo to obtain 335 mg of the the title compound as a yellowish powder.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.70 (m, 1H), 2.98 (d, *J* = 7.1 Hz, 2H), 3.57 (m, 2H), 7.00 (m, 2H), 7.5-8.00 (m, 10H), 10.66 (s, 1H), 12.80 (broad s, 1H). ESI-MS: *m/z* 461, 260, 200, 182, 115, 105. Anal. Found: C, 53.95; H, 4.65; N, 3.97; S, 18.55.

#### EXAMPLE 15

Preparation of 3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid. PNU230964




Following the general procedure outlined in Example 14 and making non-critical variations but starting with thiophenol in step 5, the title compound was obtained as a white powder.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  2.70 (m, 1H), 3.15 (m, 2H), 3.57 (dd,  $J = 4.4$  and 14.4 Hz, 1H), 3.63 (dd,  $J = 8.1$  and 14.4 Hz, 1H), 7.2-7.6 (m, 8H), 7.8-8.1 (m, 6H), 10.66 (s, 1H), 12.80 (broad s, 1H). ESI-MS:  $m/z$  455, 331, 260, 194, 123, 110, 105, 77.

Anal. Found C, 60.03; H, 5.29; N, 3.28; S, 11.68.

#### EXAMPLE 16

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-(4-hydroxyphenyl)sulfanyl]methyl]propanoic acid.




A solution of 2-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]methylacrylic acid (3.4 g) and 4-hydroxythiophenol (3 g) in 20 mL of DMF was heated for 5 h at 60°C. After cooling, the residue was chromatographed on silica gel eluting with  $\text{CHCl}_3$ /cyclohexane

1/2. The fractions containing the product were pooled and the solvent removed. The residue was crystallized from  $\text{CHCl}_3/\text{cyclohexane}$  1/3 affording 2.7 g of the title compound as a white powder.

$^1\text{H NMR}$  (400 MHz, ,  $\text{DMSO}-d_6$ ):  $\delta$  2.58 (m, 1H), 2.95 (m, 2H), 3.64 (m, 2H),  
 5 6.63 (d,  $J = 8.5$  Hz, 2H), 7.06 (d,  $J = 8.5$  Hz, 2H), 7.58 (d,  $J = 8.5$  Hz, 2H), 7.79 (d,  $J = 8.5$   
 Hz, 2H), 7.86 (d,  $J = 8.5$  Hz, 2H), 7.92 (d,  $J = 8.5$  Hz, 2H), 9.60 (broad s, 1H), 12.80 (br  
 s, 1H). MS (EI):  $m/z$  462, 251, 210, 152, 126. Anal. Found C, 56.95; H, 4.25; S,  
 13.68.

#### EXAMPLE 17

10 Preparation of 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid, sodium salt.




15 To a stirred solution of  $\alpha$ -bromomethylacrylic acid (4.95 g) and  $\text{NaHCO}_3$  (4 g) in 25 mL of DMF was added phenol (4.5 g) and the resulting suspension was heated at  $75^\circ\text{C}$  for 5 h. After disappearance of  $\alpha$ -bromomethylacrylic acid, checked by tlc and HPLC, the solution was cooled at  $-15^\circ\text{C}$  and 4'-chloro[1,1'-biphenyl]-4-thiol (9.2 g) was added. After stirring 1 h at room temperature, the solution was diluted with ethyl  
 20 acetate and washed thoroughly with 0.1 M citric acid. After drying and removal of the solvent, the residue was chromatographed on silica gel eluting with  $\text{CHCl}_3/\text{cyclohexane}$  1/2 to provide 4.9 g of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-phenoxymethyl]propanoic acid as a white foam. A solution of this intermediate (4.6 g) in 50 mL of ethanol was added dropwise to a stirred solution of Oxone<sup>®</sup> (15 g) in 100

mL of DMF at 45°C. After stirring for 4 h, the solvent was evaporated off and the residue was partitioned between chloroform and water. After drying and removal of the solvent, the crude residue was chromatographed on silica gel eluting with chloroform to provide 3.9 g of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-phenoxymethyl]propanoic acid as white foam. To a solution of this intermediate (2.15 g) in 100 mL of EtOH was added 1 M NaOH (6 mL). The resulting solution was charcoalized then filtered. Concentration in vacuum afforded 1.9 g of the title compound as shiny crystals.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.55 (m, 1H), 3.44 (dd, *J* = 8.3 and 14.6 Hz, 1H), 3.97 (dd, *J* = 3.7 and 14.6 Hz, 1H), 4.01 (dd, *J* = 4.4 and 8.8 Hz, 1H), 4.20 (dd, *J* = 4.4 and 8.8 Hz, 1H), 6.75 (d, *J* = 8.8 Hz, 2H), 6.83 (t, *J* = 7.3 Hz, 1H), 7.18 (dd, *J* = 7.3 and 8.8 Hz, 2H), 7.56 (d, *J* = 8.8 Hz, 2H), 7.72 (d, *J* = 8.8 Hz, 2H), 7.85 (d, *J* = 8.5 Hz, 2H), 7.91 (d, *J* = 8.5 Hz, 2H). ESI(-)MS: *m/z* 429, 291, 251. Anal. Found C, 55.54; H, 4.52; S, 6.74.

#### EXAMPLE 18

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid.



To stirred solution of glycine ethyl ester hydrochloride (6. g) and TEA (16 mL) in 80 mL of DMF was added α-bromomethylacrylic acid (5 g) at 0°C. After stirring for 2 h at rt, the suspension was filtered and the filtrate was evaporated to dryness. The residue dissolved in 50 mL of dioxane was treated with *n*-butyl isocyanide (4 mL) and

heated for 5 h at 65°C. The solvent was removed and the crude residue was columned over a small pad of silica gel eluting with dichloromethane to afford 5.5 g of 2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)-methyl]acrylic acid as an oil. This intermediate (5.5 g) was dissolved in DMF (50 mL) and treated at 0 °C with 4'-chloro[1,1'-biphenyl]-4-thiol (5.5 g) and NaHCO<sub>3</sub> (3.9 g). After stirring 1 h at rt, the reaction mixture was concentrated in vacuo, taken up in ethyl acetate, and sequentially washed with 0.5 M HCl and brine. After drying over anhydrous sodium sulfate, the solvent was removed and the residue was chromatographed on silica eluting with ethyl acetate/cyclohexane 1/1 to provide 4.6 g of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfanyl]-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid. A solution of this intermediate (4.5 g) and Oxone<sup>®</sup> (13 g) in 100 mL of DMF was heated for 1 h at 50°. The reaction mixture was concentrated in vacuo and the residue was partitioned between ethyl acetate and water. After drying over sodium sulfate, the solvent was removed in vacuo and the residue was columned over a small pad of silica gel eluting with CH<sub>2</sub>Cl<sub>2</sub> to give the title compound (4.2 g) as a white powder.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 0.83 (t, *J* = 7.3 Hz, 3H), 1.20 (m, 2H), 1.40 (m, 2H), 3.00 (m, 1H), 3.30 (m, 2H), 3.52 (d, *J* = 7.6 Hz, 2H), 3.53 (m, 1H), 3.71 (dd, *J* = 8.3 and 14.9 Hz, 1H), 3.79 and 3.93 (two d, *J* = 17.3 Hz, 2H), 7.58 (d, *J* = 8.5 Hz, 2H), 7.79 (d, *J* = 8.5 Hz, 2H), 7.94 (s, 4H), 12.90 (broad s, 1H). ESI(+)MS: *m/z* 493, 475, 319, 235, 223. Anal. Found C, 55.38; H, 5.19; N, 5.33; S, 6.78.

#### EXAMPLE 19

Preparation of 3-[(4'-cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid.




By following the procedure described in Example 1, steps 4-6, and making non-critical variations, but replacing 4'-chloro[1,1'-biphenyl]-4-thiol with 4'-cyano[1,1'-biphenyl]-4-thiol, the title compound was obtained as a white powder.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 3.09 (m, 2H), 3.36 (m, 1H), 3.61 (dd, *J* = 3.8 and 14.5 Hz, 1H), 3.70 (dd, *J* = 7.7 and 14.5 Hz, 1H), 7.20 (m, 5H), 7.6-7.7 (m, 6H), 7.94 (d, *J* = 8.5 Hz, 2H). Anal. Found C, 62.83; H, 4.41; N, 3.13; S, 14.29.

#### EXAMPLE 20

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]-propanoic acid.




Following the general procedure outlined in Example 16 and making non-critical variations but replacing 4-hydroxy-thiophenol with 2-mercaptopyridine, the title compound was obtained as a white crystalline powder.

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 2.99 (m, 1H), 3.37 (d, *J* = 7.5 Hz, 1H), 3.66 (dd, *J* = 4.1 and 14.6 Hz, 1H), 3.70 (dd, *J* = 8.3 and 14.6 Hz, 1H), 7.01 (ddd, *J* = 1.0, 4.9 and 7.4 Hz, 1H), 7.18 (ddd, *J* = 0.9, 1.0 and 8.1 Hz, 1H), 7.50 (ddd, *J* = 1.9, 7.4 and 8.1 Hz, 1H), 7.56 (d, *J* = 8.6 Hz, 2H), 7.76 (d, *J* = 8.6 Hz, 2H), 7.88 (m, 4H), 8.30 (ddd, *J* = 0.9, 1.9 and 4.9 Hz, 1H), 12.85\* (broad s, 1H). ESI(+)MS: *m/z* 448, 319, 235, 178. Anal. Found C, 56.26; H, 4.09; N, 3.17; S, 14.30.

#### EXAMPLE 21

Preparation of 3-[[4-[(4-chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]-propanoic acid.






Following the general procedure outlined in Example 14 and making non-critical  
 5 variations but starting with 4-chlorobenzoyl chloride in step 1, and replacing 2-  
 thiophenethiol with thiophenol in step 5, the title compound was obtained as a white  
 powder.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.67 (m, 1H), 3.15 (m, 2H), 3.62 (m, 2H),  
 7.22 (m, 5H), 7.64 (d, *J* = 8.5 Hz, 2H), 7.77 (d, *J* = 8.0 Hz, 2H), 8.01 (m, 4H), 10.70 (s,  
 10 1H), 12.80 (broad s, 1H). ESI(-)MS: *m/z* 510 [M + Na - 2H]<sup>+</sup>, 334. Anal. Found C,  
 55.66; H, 4.10; N, 2.91; S, 12.82.

#### EXAMPLE 22

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-  
 pyrimidinylsulfanyl)methyl]-propanoic acid.



Step 1. 3-Bromo-2-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]propanoic acid.

2-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]methyl]acrylic acid (4 g), prepared  
 as described in Example 1, step 5, was suspended in 33% HBr in acetic acid (40 mL).  
 20 After stirring for 4 h at ambient temperature, the reaction mixture was concentrated in

vacuo, taken up in toluene and evaporated to dryness (twice). The residue was triturated with ethyl ether, whereby the title product separated as a white crystalline powder, which was collected by filtration (3.85 g). A second crop (0.3 g) was recovered from the mother liquors by repeating the process after evaporation to dryness.


- 5 Step 2. 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfanyl)methyl]propanoic acid.

2-Mercaptopyrimidine (3.65 g) was added portionwise to a suspension of sodium hydride (60% dispersion in mineral oil; 1.21 g) in anhydrous DMF (80 mL) under an argon blanket, and the mixture stirred for 20 min at room temperature to obtain  
 10 a clear solution, after which time the bromomethylpropanoic acid from step 1 above (4.54 g) was added. After 4 h at room temperature, 0.2 N HCl (600 mL) was added, and the mixture was extracted twice with ethyl acetate. The organic extract was washed with brine, dried over sodium sulfate, evaporated, taken up in toluene and evaporated again to dryness. Trituration of the residue with ethyl ether afforded a powder (2.99 g), which  
 15 was purified by stirring for 1 h in a mixture of ether/ethyl acetate 9/1 and filtration, thereby obtaining the title product (2.7 g) as a white powder, mp. 166-168°C.

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 3.06 (m, 1H), 3.28 (partially overlapped by water, 2H), 3.62 (dd, *J* = 4.0 and 14.6 Hz, 1H), 3.72 (dd, *J* = 8.2 and 14.6 Hz, 1H), 7.10 (t, *J* = 4.9 Hz, 1H), 7.57 (d, *J* = 8.5 Hz, 2H), 7.75 (d, *J* = 8.5 Hz, 2H), 7.90 (m, 4H), 8.50  
 20 (d, *J* = 4.9 Hz, 2H), 13.00 (broad s, 1H). ESI(+)MS: *m/z* 449, 319, 235, 179, 113. Anal. Found C, 53.65; H, 3.92; N, 6.21; S, 14.30.

### EXAMPLE 23

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-amino-1,3,4-thiadiazol-2-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 5-amino-2-mercapto-1,3,4-thiadiazole, the title compound was obtained as a yellowish powder, mp 184-186 °C.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.98 (m, 1H), 3.30 (m, 2H), 3.68 (m, 2H), 7.33 (s, 2H), 7.57 (d, *J* = 8.5 Hz, 2H), 7.78 (d, *J* = 8.5 Hz, 2H), 7.92 (m, 4H), 13.00 (broad s, 1H). ESI(+)-MS: *m/z* 470, 337, 319, 235. Anal. Found C, 41.38; H, 3.42; N, 10.83; S, 22.89.

#### EXAMPLE 24

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-methyl-2H-1,2,3-triazol-4-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 4-mercapto-2-methyl-2H-1,2,3-triazole, the title compound was obtained as a white powder, mp 166-168 °C.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 3.02 (m, 1H), 3.34 (m, 2H), 3.48 (m, 3H), 3.69 (m, 2H), 7.58 (d, *J* = 8.5 Hz, 2H), 7.79 (d, *J* = 8.5 Hz, 2H), 7.91 (m, 4H), 8.49 (s,

1H), 12.98 (broad s, 1H). ESI(+)MS: m/z 452, 337, 319, 235. Anal. Found: C, 50.18; H, 4.06; N, 9.25; S, 13.45.

#### EXAMPLE 25

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[1-methyl-1H-imidazol-2-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2-mercapto-1-methyl-1H-imidazole, the title compound was obtained as a white powder.

<sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>): δ 2.90 (m, 1H), 3.20 (m, 2H), 3.63 (m, 2H), 6.88 (d, J = 2.1 Hz, 1H), 7.16 (d, J = 2.1 Hz, 1H), 7.56 (d, J = 8.5 Hz, 2H), 7.78 (d, J = 8.5 Hz, 2H), 7.91 (m, 4H). ESI(+)MS: m/z 451, 337, 235, 199, 181.

#### EXAMPLE 26

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2,6-dimethyl-4-pyrimidinyl)-sulfanyl]methyl]propanoic acid.



Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2,6-dimethyl-4-


mercaptopyrimidine, the title compound was obtained as a white powder, mp 170-172 °C.

ESI(+)-MS: m/z 477, 336, 251, 187, 152, 140. Anal. Found C, 55.05; H, 4.52; N, 4.64; S, 12.32.

5

#### EXAMPLE 27

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4,6-dimethyl-2-pyrimidinyl)-sulfanyl]methyl]propanoic acid.



10

Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 4,6-dimethyl-2-mercaptopyrimidine, the title compound was obtained as a white powder, mp 170-172 °C.


<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 2.20 (s, 6H), 3.05 (m, 1H), 3.30 (m, 2H), 3.61 (dd, *J* = 3.4 and 14.5 Hz, 1H), 3.73 (d, *J* = 8.1 and 14.5 Hz, 1H), 6.81 (s, 1H), 7.58 (d, *J* = 8.5 Hz, 2H), 7.75 (d, *J* = 8.5 Hz, 2H), 7.88 (m, 4H), 12.90 (broad s, 1H). ESI(+)-MS: m/z 477, 319, 294, 235, 225, 141. Anal. Found C, 53.90; H, 4.52; N, 5.70; S, 12.48.

15

#### EXAMPLE 28

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1,3-thiazol-2-ylsulfanyl)-methyl]propanoic acid.

20




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2-mercaptothiazole, the title compound was obtained as a white powder, mp 152-154 °C.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 3.00 (m, 1H), 3.45 (m, 2H), 3.63 (dd, *J* = 4.7 and 14.5 Hz, 1H), 3.74 (d, *J* = 7.7 and 14.5 Hz, 1H), 7.57 (m, 3H), 7.65 (d, *J* = 3.4 Hz, 1H), 7.60 (d, *J* = 8.5 Hz, 2H), 7.77 (d, *J* = 8.5 Hz, 2H), 7.91 (m, 4H), 13.00 (broad s, 1H). ESI(+)MS: *m/z* 454, 337, 319, 294, 235, 184. Anal. Found: C, 50.07; H, 3.69; N, 3.15; S, 21.14.

#### EXAMPLE 29

Preparation of 3-[(4'-cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxy)methylpropanoic acid, sodium salt.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing 4'-chloro[1,1'-biphenyl]-4-thiol with 4'-cyano[1,1'-biphenyl]-4-thiol, the title compound was obtained as a white powder.

Anal. Found C, 59.59; H, 4.45; N, 2.88; S, 6.85.

#### EXAMPLE 30

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-acetylaminophenoxy)-

methyl]propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing phenol with 4-acetamidophenol, the title compound was obtained as a white powder.

Anal. Found C, 58.03; H, 4.61; N, 2.75; S, 6.84.

#### EXAMPLE 31


Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[4-(4-hydroxy-6-methyl-2-pyrimidinyl)sulfanyl]methyl]propanoic acid.



Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 4-hydroxy-6-methyl-2-mercaptopyrimidine, the title compound was obtained as a white powder, mp 165-168 °C.

#### EXAMPLE 32

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1-methyl-1H-tetrazol-5-yl]-sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 1-methyl-1H-tetrazole-5-thiol, the title compound was obtained as a whitish powder, mp 98-100 °C.

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 3.08 (m, 1H), 3.54 (m, 2H), 3.70 (dd, *J* = 4.3 and 14.6 Hz, 1H), 3.76 (dd, *J* = 7.6 and 14.6 Hz, 1H), 3.85 (s, 3H), 7.59 (d, *J* = 8.5 Hz, 2H), 7.79 (d, *J* = 8.5 Hz, 2H), 7.92 (m, 4H), 13.00 (broad s, 1H). ESI (+) MS: *m/z* 453, 337, 319, 294, 235. Anal. Found C, 47.64; H, 4.59; N, 13.18; S, 11.86.

### EXAMPLE 33

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(hydroxymethyl)phenoxy]-methyl]propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing phenol with salicyl alcohol, the title compound was obtained as a crystalline powder.

Anal. Found C, 59.50; H, 4.62; S, 6.98.



## EXAMPLE 34

Preparation of 3-[[4-[(4-chlorobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxyethyl)propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing 4-chloro[1,1'-biphenyl]-4-thiol with 4-chloro-N-(4-sulfanylbenzamide, the title compound was obtained as a white powder.

Anal. Found C, 57.72; H, 4.28; N, 3.02; S, 6.62.

## EXAMPLE 35

Preparation of 3-[[4-[(4-cyanobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxyethyl)propanoic acid.




Following the general procedure outlined in Example 17 and making non-critical variations but replacing 4-chloro[1,1'-biphenyl]-4-thiol with 4-cyano-N-(4-sulfanylbenzamide, the title compound was obtained as a white powder.

Anal. Found C, 61.33; H, 4.26; N, 6.19; S, 7.15.

## EXAMPLE 36

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]methyl]propanoic acid.




Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 5-methyl-2-mercapto-1,3,4-thiadiazole, the title compound was obtained as a withish powder, mp 147-150 °C.

Anal. Found, C, 47.03; H, 3.74; N, 5.69; S, 11.87.

## EXAMPLE 37

Preparation of 3-[(4'-chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1H-benzimidazol-2-yl)sulfanyl]methyl]-propanoic acid.



Following the general procedure outlined in Example 22 and making non-critical variations but replacing 2-mercaptopyrimidine with 2-mercapto-1H-benzimidazole, the title compound was obtained as a withish powder, mp 176-179 °C.

Anal. Found C, 55.43; H, 3.96; N, 5.57; S, 12.32.

## EXAMPLE 38

### Biochemical Activity Test

The inhibitory activity of the compounds of the present invention was evaluated in vitro against one or more of the MMP enzymes (gelatinases, collagenases, stromelysins) using the internal fluorescence quenching method.. The assay method is based on the hydrolysis of the MMP substrate, MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH<sub>2</sub> at 37 °C (see Knight, C.G. et al., FEBS Lett. 296:263-266, 1992). The enzymes cleave at the Gly-Leu bond removing the internally quenching DPA group. The release of the highly fluorescent peptide MCA-Pro-Leu was monitored with a LS50 Perkin Elmer fluorimeter equipped with a Peltier-thermostated cell holder using an excitation wavelength of 326 nm and an emission wavelength of 398 nm. Slits were set at 14/14. Proper determination of steady-state rates of substrate cleavage required a preincubation at 37 °C of 15 minutes to allow for complete equilibration of the enzyme-inhibitor complex. For the determination of the apparent  $K_i$  ( $K_{i,app}$ ), the concentration of the inhibitor was varied at a constant and low concentration of the substrate (3  $\mu$ M). Rates of substrate hydrolysis, expressed as fluorescence arbitrary units per minute, were calculated by least-square analysis of the steady-state linear progression curves. The steady-state rates were fitted to the Michaelis equation describing competitive inhibition; in particular,  $K_{i,app}$  were determined by fitting the data to the tight-binding equation of Morrison (Biochem. Biophys. Acta, 185:269-286, 1969) by non-linear methods. Since the Michaelis constants ( $K_m$ ) for the MCA peptide substrate with the MMPs is quite high (70  $\mu$ M or greater; see Knight, C.G. et al., reference above) and by far exceeds the substrate concentration, the factor  $(1 + S/K_m)$  can be approximated to unity without appreciable error, so that the determined  $K_{i,app}$  will be essentially equal to  $K_i$ .

Human gelatinase-A (MMP-2) was obtained as pro-enzyme (72 kDa) and was activated with 1 mM 4-aminophenylmercuric acetate for 30 min at 37 °C immediately prior to use. The inhibitor was diluted into the assay from a stock solution in 100% DMSO, and controls substituted an equal volume of DMSO so that the final DMSO concentration from inhibitor and substrate dilution in all assays was 1%. The

concentration of MMP-2 was 19.2 pM. For compounds of formula (I) where X is -OH, the assay buffer was 50 mM Tris/HCl, pH 6.5, containing 150 mM NaCl, 10 mM CaCl<sub>2</sub>, 0.01 mM ZnCl<sub>2</sub> and 0.05% Brij-35 (Protocol A). Other conditions were as stated above.

Human fibroblast collagenase (MMP-1) was obtained as truncated recombinant enzyme encompassing residues 101-269 and did not require activation. Assay conditions were the same as above, but enzyme concentration was 200 pM.

Inhibitory data (K<sub>i</sub>'s against MMP-2 and MMP-1, all nM; Protocol A) for some representative compounds of formula (I) wherein X is -OH are displayed in Table

1.

TABLE 1

Example No.	MMP-2 K <sub>i</sub> (nM)	MMP-1 K <sub>i</sub> (nM)
1	8.3	> 2500
3, Isomer 1	6.0	6600
3, Isomer 2	660	33190
5	5.5	> 2000
6, Isomer 1	3.7	3600
6, Isomer 2	50	n.t.
7	42	n.t.
9	3	n.t.
11	4	n.t.
13	12	5286
14	8	n.t.
15	7	81090
16	9	9697
17	9.2	14740

	18	7.8	n.t.
	19	17	n.t.
	20	3.9	6291
	21	1.4	50560
5	22	8	8033
	23	25	n.t.
	24	24	n.t.
	25	16	n.t.
10	26	26.8	n.t.
	27	22.8	n.t.
	28	6.6	n.t.
	29	21.5	n.t.
	30	46	19580
15	32	26	n.t.
	33	15	n.t.
	34	2.4	53980
	35	4.8	67520
	36	33.4	n.t.
20	37	18.2	n.t.

The compounds of formula (I) where X is -NHOH were tested against MMP-2 and MMP-1 under the conditions stated above, but the assay solution was buffered at pH 7.4 (Protocol B). Inhibitory data (K<sub>i</sub>'s, all picoM) for some representative compounds of formula (I) wherein X is -NHOH are displayed in Table 2.

TABLE 2


Example No.	MMP-2 K <sub>i</sub> (picoM)	MMP-1 K <sub>i</sub> (picoM)
4	159	149000

8	107	26800
10	50 <del>2</del>	12600 <del>2</del>
12	69 <del>2</del>	17100 <del>2</del>

---

# CLAIMS

1. A compound which is a 3-arylsulfonyl-2-methyl propanoic acid derivative of formula (I):



wherein

X is HO-NH- or HO-;

10  $R_1$  is selected from phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-cyanophenyl, benzamido (i.e., -NH-CO-Ph) and benzamido substituted on the terminal phenyl ring by  $C_1$ - $C_4$  alkyl, fluoro, chloro, cyano or  $C_1$ - $C_4$  alkoxy;

$R_2$  is selected from:

(a) -S-Ar or -S-CH<sub>2</sub>-Ar, wherein Ar is a monocarbocyclic or  
15 bicarbocyclic aromatic moiety which is either unsubstituted or substituted with one or two substituents selected from  $C_1$ - $C_4$  alkyl, phenyl, benzyl,  $C_1$ - $C_4$  alkoxy, fluoro, chloro, bromo, nitro, cyano, hydroxy, amino, dimethylamino, acetamido, methylthio and acetyl;

(b) -O-Ar, wherein Ar is as defined above;


(c) -S-Het or -S-CH<sub>2</sub>-Het, wherein Het is a heterocyclic ring selected  
20 from pyridine, pyrimidine, pyridazine, pyrazine, 1,2,5-triazine, imidazole, thiophene, furan, pyrrole, pyrazole, 1,3-thiazole, 1,3-oxazole, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole, 1,2,3,4-tetrazole, quinoline, isoquinoline, indole, 1,3-benzoxazole, 1,3-benzothiazole, benzimidazole, [1,3]oxazolo[4,5-b]pyridine, [1,3]thiazolo[4,5-b]pyridine, [1,2,3,4]tetrazolo[1,5-b]pyridazine and purine, and

wherein said Het group can be substituted with one to three substituents selected from C<sub>1</sub>-C<sub>4</sub> alkyl, phenyl, pyridyl, benzyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, methylthio, fluoro, chloro, nitro, cyano, hydroxy, oxo, amino, methylamino, dimethylamino, 2-dimethylaminoethyl, acetamido and acetyl; and

(d) 2,5-dioxo-1-imidazolidinyl or 2,4-dioxo-1-imidazolidinyl, either of which is optionally substituted at the carbon atom by one or two methyl, linear or branched C<sub>2</sub>-C<sub>4</sub> alkyl, phenyl, benzyl or hydroxymethyl groups, and at the nitrogen atom with C<sub>1</sub>-C<sub>4</sub> linear or branched alkyl;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, which is an isomer having the configuration depicted in formula (I):



wherein X, R<sub>1</sub> and R<sub>2</sub> are as defined in claim 1.

3. A compound according to claim 1 which is selected from:

- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;
- (2S)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfonyl)methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfonyl)methyl]-propanamide;



3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-hydroxyphenyl)sulfanyl]methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4-hydroxyphenyl)sulfanyl]-methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylsulfanyl)methyl]-propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylmethyl)sulfanyl]methyl]-propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2-pyridinylmethyl)sulfanyl]-methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(5-methyl-1,3,4-thiadiazol-2-yl)sulfanyl]methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-amino-1,3,4-thiadiazol-2-yl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-methyl-1H-imidazol-2-yl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyrimidinylsulfonyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-methyl-1H-tetrazol-5-yl)sulfonyl]methyl]-

5 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1-benzyl-1H-tetrazol-5-yl)sulfonyl]methyl]-

propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1H-benzimidazol-2-ylsulfonyl)methyl]-

10 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(1,3-thiazol-2-ylsulfonyl)methyl]propanoic

acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-methyl-2H-1,2,3-triazol-4-yl)sulfonyl]-

15

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-phenyl-1,3,4-oxadiazol-2-yl)sulfonyl]-

methyl]propanoic acid;

20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1-(3-pyridinyl)-1H-imidazol-2-yl]sulfonyl]-

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-methyl-1,3-oxazol-2-yl)sulfonyl]methyl]-

25 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1,3]oxazolo[4,5-b]pyrimidin-2-yl]sulfonyl]-

methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[1,2,3,4]tetrazolo[1,5-b]pyridazin-6-yl]sulfonyl]methyl]propanoic acid;

30

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(2,6-dimethyl-4-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4,6-dimethyl-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

5 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4-hydroxy-6-methyl-2-pyrimidinyl)-sulfanyl]methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[[(4,6-dihydroxy-2-pyrimidinyl)sulfanyl]-methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(9H-purin-6-ylsulfanyl)methyl]propanoic  
10 acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylaminophenoxy)methyl]propanoic

15 acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4-acetylaminophenoxy)methyl]propanoic

acid;

3-[(1,1'-Biphenyl)-4-ylsulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]-  
20 propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-

methyl]propanoic acid;

25 (2*R*)-3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanoic acid;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-*N*-hydroxy-2-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;

3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)-  
30

- methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(4,4-dimethyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2,5-dioxo-1-imidazolidinyl)methyl]propanoic
- 5 acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(2,5-dioxo-1-imidazolidinyl)-
- methyl]propanamide;
- 10 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-methyl-2,5-dioxo-1-imidazolidinyl)-
- methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-methyl-2,5-dioxo-1-imidazolidinyl)methyl]-
- propanoic acid;
- 15 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]-
- propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;
- 20 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]-
- propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,4-dioxo-1-imidazolidinyl)methyl]propanamide;
- 25 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-isopropyl-2,4-dioxo-1-imidazolidinyl)-
- methyl]propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;

- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-3-methyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 5 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Fluoro[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 10 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-(phenoxymethyl)propanamide;
- 15 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-acetylamino)phenoxymethyl]propanoic acid;
- 20 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylsulfanyl)methyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(2-pyridinylmethyl)sulfanyl)methyl]propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]-
- 25 propanoic acid;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-N-hydroxy-2-[(3-butyl-2,5-dioxo-1-imidazolidinyl)methyl]propanamide;
- 3-[(4'-Cyano[1,1'-biphenyl]-4-yl)sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-imidazolidinyl)methyl]propanoic acid;
- 30

3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;

3-[[4-(Benzoylamino)phenyl]sulfonyl]-N-hydroxy-2-[(phenylsulfanyl)methyl]-  
propanamide;

3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[[2-

5 pyridinylmethyl)sulfanyl)methyl]propanoic  
acid;

3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-[(2-thienylsulfanyl)methyl]propanoic acid;

3-[[4-(Benzoylamino)phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic  
10 acid;

3-[[4-[(4-Methylbenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-  
[(phenylsulfanyl)methyl]-  
propanamide;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic  
15 acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-  
[(phenylsulfanyl)methyl]-  
propanamide;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;

20 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-  
propanamide;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic  
acid;


3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[[2-  
25 pyridinylmethyl)sulfanyl)methyl]-  
propanoic acid;

3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-  
yl)sulfanyl]-  
methyl]propanoic acid;

30 3-[[4-[(4-Chlorobenzoyl)amino]phenyl]sulfonyl]-2-[(5-hydroxymethyl-2,4-dioxo-1-

- imidazolidinyl)methyl]propanoic acid;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(phenylsulfanyl)methyl]propanoic acid;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-  
 5 [(phenylsulfanyl)methyl]-  
 propanamide;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-  
 methyl]propanoic acid;
- 10 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(phenoxymethyl)-  
 propanamide;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-(3-fluorophenoxymethyl)propanoic acid;
- 15 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-N-hydroxy-2-(3-fluorophenoxymethyl)-  
 propanamide;
- 3-[[4-[(4-Cyanobenzoyl)amino]phenyl]sulfonyl]-2-[(2-acetyl amino)phenoxymethyl]-  
 propanoic acid;
- 20 3-[[4-(4-Propylbenzoylamino)phenyl]sulfonyl]-2-[[1-methyl-1H-tetrazol-5-yl)sulfanyl]-  
 methyl]propanoic acid; and
- 3-[[4-[(4-Methoxybenzoyl)amino]phenyl]sulfonyl]-2-(phenoxymethyl)propanoic acid;
- 3-[(4'-Chloro[1,1'-biphenyl]-4-yl)sulfonyl]-2-[[2-(hydroxymethyl)phenoxy]methyl]-  
 25 propanoic acid; and the pharmaceutically acceptable salts thereof.

4. A process for producing a compound as defined in claim 1, starting from a compound of formula 4:



wherein R is H or the residue of a carboxylic acid ester,  $R_1$  and  $R_2$  are as defined in claim 1 and n is 0 or 2, which process comprises:

(a) hydrolysing a said compound of formula 4 in which R is the residue of a carboxylic and ester to give a compound of formula (I) in which X is HO-; or

(b) hydrolysing and oxidising, in either order, a said compound of formula 4 in which n is 0 and R is the residue of a carboxylic acid ester, to give a compound of formula (I) in which X is HO-; or

(c) activating a said compound of formula 4 wherein R is H and n is 2 to form an activated carboxy group, coupling the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotecting the hydroxamic group to give a compound of formula (I) wherein X is -NHOH; or


(d) submitting a said compound of formula 4 wherein R is H and n is zero to a sequence of reactions comprising oxidation at the sulphur atom, activation of the carboxy group, condensation of the activated carboxy group with hydroxylamine or an O-protected derivative thereof and, if necessary, deprotection of the hydroxamic group to form a compound of formula (I) wherein X is -NHOH, the oxidation step being conducted either before the activation step or after the condensation step; and/or

(e) if desired, converting a resulting compound of formula (I) into another compound of formula (I) and/or converting a free compound into a pharmaceutically acceptable salt thereof, and/or converting a salt into a free compound.

5. A process according to claim 4 wherein the compound of formula 4 is obtained by


(a) subjecting a compound of formula 2:






wherein R, R<sub>1</sub> and n are as defined in claim 4, to conjugate addition by treatment with a compound of formula R<sub>2</sub>H wherein R<sub>2</sub> is as defined in claim 1; or

(b) treating a compound of formula 3:



wherein R and R<sub>2</sub> are as defined in claim 4, with a thiol of formula:



to obtain a compound of formula 4 in which n is zero.

6. A pharmaceutical composition which comprises a pharmaceutically acceptable carrier or diluent and, as an active principle, a compound as defined in any one of claims 1 to 4.

7. A compound as defined in any one of claims 1 to 4, for use in a method of treatment of the human or animal body by therapy.


8. A compound as claimed in claim 7 for use in the treatment or prophylaxis of a disease in a mammal mediated by a matrix metalloproteinase.

9. A compound as claimed in claim 8 wherein the matrix metalloproteinase is a gelatinase (MMP-2), the membrane-type-MMPs involved in gelatinase activation (MMP-14), a stromelysin (MMP-3 or MMP-10), collagenase (MMP-13) or neutrophil collagenase (MMP-8).

10. A compound as claimed in any one of claims 7 to 9 for use in the treatment or prophylaxis of tumour growth or metastasis, rheumatoid arthritis, osteoarthritis, ophthalmic disease, cardiovascular disease, periodontal disease, multiple sclerosis or Alzheimer's disease.

ABSTRACT**3-ARYLSULFONYL-2-(SUBSTITUTED-METHYL) PROPANOIC ACID  
DERIVATIVES AS MATRIX METALLOPROTEINASE INHIBITORS**

Compounds which are 3-arylsulfonyl-2-methyl propanoic acid derivatives of formula (I):



wherein

X is HO-NH- or HO-

R<sub>1</sub> is selected from phenyl, 4-chlorophenyl, 4-fluorophenyl, 4-cyanophenyl, benzamido (i.e., -NH-CO-Ph) and benzamido substituted on the terminal phenyl ring by C<sub>1</sub>-C<sub>4</sub> alkyl, fluoro, chloro, cyano or C<sub>1-4</sub> alkoxy;

R<sub>2</sub> is selected from

- (a) -S-Ar or -S-CH<sub>2</sub>-Ar wherein Ar is an aromatic moiety;
- (b) -O-Ar wherein Ar is as defined above;
- (c) -S-Het or -S-CH<sub>2</sub>-Het wherein Het is a heterocyclic ring; and
- (d) 2,5-dioxo-1-imidazolidinyl or 2,4-dioxo-1-imidazolinyl; and the

pharmaceutically acceptable salts thereof; have potent and selective inhibitory activity against matrix metalloproteinases (MMPs) and can thus be used in the treatment and prevention of diseases mediated by MMPs.

**THIS PAGE BLANK (USPTO)**